

Title	Strong Convergence Theorem by the Hybrid and Extragradient Method for Nonexpansive Mappings and Monotone Mappings (Advanced Topics of Information Science and Functional Analysis)
Author(s)	Nadezhkina, Natalia; Takahashi, Wataru
Citation	数理解析研究所講究録 (2004), 1396: 42-48
Issue Date	2004-10
URL	http://hdl.handle.net/2433/25965
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Strong Convergence Theorem by the Hybrid and Extragradient Method for Nonexpansive Mappings and Monotone Mappings

Natalia Nadezhkina and Wataru Takahashi

Department of Mathematical and Computing Sciences
Graduate School of Information Science and Engineering
Tokyo Institute of Technology

Abstract

In this paper we introduce an iterative process for finding a common element of the set of fixed points of a nonexpansive mapping and the set of solutions of a variational inequality problem for a monotone, Lipschitz continuous mapping. The iterative process is based on two well known methods - hybrid and extragradient. We obtain a strong convergence theorem for three sequences generated by this process.

1 Introduction

Let C be a closed convex subset of a real Hilbert space H and let P_C be the metric projection of H onto C . A mapping A of C into H is called *monotone* if

$$\langle Au - Av, u - v \rangle \geq 0$$

for all $u, v \in C$. The *variational inequality problem* is to find a $u \in C$ such that

$$\langle Au, v - u \rangle \geq 0$$

for all $v \in C$. The set of solutions of the variational inequality problem is denoted by $VI(C, A)$. A mapping A of C into H is called α -*inverse-strongly-monotone* if there exists a positive real number α such that

$$\langle Au - Av, u - v \rangle \geq \alpha \|Au - Av\|^2$$

for all $u, v \in C$; see [1], [4]. It is obvious that an α -inverse-strongly-monotone mapping A is monotone and Lipschitz-continuous. A mapping S of C into itself is called *nonexpansive* if

$$\|Su - Sv\| \leq \|u - v\|$$

for all $u, v \in C$; see [8]. We denote by $F(S)$ the set of fixed points of S . For finding an element of $VI(C, A)$ under the assumption that a set $C \subset H$ is closed and convex and a mapping A of C into H is α -inverse-strongly-monotone, Iiduka, Takahashi and Toyoda [2] introduced the following iterative scheme by a hybrid method:

$$\begin{cases} x_0 = x \in C \\ y_n = P_C(x_n - \lambda_n A x_n) \\ C_n = \{z \in C : \|y_n - z\| \leq \|x_n - z\|\} \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\} \\ x_{n+1} = P_{C_n \cap Q_n} x \end{cases}$$

for every $n = 0, 1, 2, \dots$, where $\lambda_n \subset [a, b]$ for some $a, b \in (0, 2\alpha)$. They showed that if $VI(C, A)$ is nonempty, then the sequence $\{x_n\}$, generated by this iterative process, converges strongly to $P_{VI(C, A)}x$. On the other hand, for solving the variational inequality problem in a finite-dimensional Euclidean space \mathbb{R}^n under the assumption that a set $C \subset \mathbb{R}^n$ is closed and convex and a mapping A of C into \mathbb{R}^n is monotone and k -Lipschitz-continuous, Korpelevich [3] introduced the following so-called extragradient method:

$$\begin{cases} x_0 = x \in R^n \\ \bar{x}_n = P_C(x_n - \lambda A x_n) \\ x_{n+1} = P_C(x_n - \lambda A \bar{x}_n) \end{cases} \quad (1)$$

for every $n = 0, 1, 2, \dots$, where $\lambda \in (0, 1/k)$. He showed that if $VI(C, A)$ is nonempty, then the sequences $\{x_n\}$ and $\{\bar{x}_n\}$, generated by (1), converge to the same point $z \in VI(C, A)$.

In this paper, by an idea of combining hybrid and extragradient methods, we introduce an iterative process for finding a common element of the set of fixed points of a nonexpansive mapping and the set of solutions of a variational inequality problem for a monotone, Lipschitz continuous mapping in a real Hilbert space. Then we obtain a strong convergence theorem for three sequences generated by this process.

2 Preliminaries

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$ and let C be a closed convex subset of H . We write $x_n \rightharpoonup x$ to indicate that the sequence $\{x_n\}$ converges weakly to x and $x_n \rightarrow x$ to indicate that $\{x_n\}$ converges strongly to x . For every point $x \in H$ there exists a unique nearest point in C , denoted by $P_C x$, such that $\|x - P_C x\| \leq \|x - y\|$ for all $y \in C$. P_C is called the *metric projection* of H onto C . We know that P_C is a nonexpansive mapping of H onto C . It is also known that P_C is characterized by the following properties: $P_C x \in C$ and

$$\langle x - P_C x, P_C x - y \rangle \geq 0; \quad (2)$$

$$\|x - y\|^2 \geq \|x - P_C x\|^2 + \|y - P_C x\|^2 \quad (3)$$

for all $x \in H, y \in C$; see [8] for more details. Let A be a monotone mapping of C into H . In the context of variational inequality problem this implies

$$u \in VI(C, A) \Leftrightarrow u = P_C(u - \lambda A u), \quad \forall \lambda > 0.$$

It is also known that H satisfies Opial's condition [6], i.e., for any sequence $\{x_n\}$ with $x_n \rightharpoonup x$ the inequality

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\|$$

holds for every $y \in H$ with $y \neq x$.

A set-valued mapping $T : H \rightarrow 2^H$ is called *monotone* if for all $x, y \in H, f \in Tx$ and $g \in Ty$ imply $\langle x - y, f - g \rangle \geq 0$. A monotone mapping $T : H \rightarrow 2^H$ is *maximal* if its graph $G(T)$ is not properly contained in the graph of any other monotone mapping. It is known that a monotone mapping T is maximal if and only if for $(x, f) \in H \times H, \langle x - y, f - g \rangle \geq 0$ for every $(y, g) \in G(T)$ implies $f \in Tx$. Let A be a monotone, k -Lipschitz-continuous mapping of C into H and $N_C v$ be the normal cone to C at $v \in C$, i.e. $N_C v = \{w \in H : \langle v - u, w \rangle \geq 0, \forall u \in C\}$. Define

$$Tv = \begin{cases} Av + N_C v, & \text{if } v \in C, \\ \emptyset, & \text{if } v \notin C. \end{cases}$$

Then T is maximal monotone and $0 \in Tv$ if and only if $v \in VI(C, A)$; see [7].

3 Strong Convergence Theorem

In this section we prove a strong convergence theorem by a combined hybrid-extragradient method for nonexpansive mappings and monotone, k -Lipshitz-continuous mappings.

Theorem 3.1 *Let C be a closed convex subset of a real Hilbert space H . Let A be a monotone and k -Lipschitz-continuous mapping of C into H and S be a nonexpansive mapping of C into itself such that $F(S) \cap VI(C, A) \neq \emptyset$. Let $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ be sequences generated by*

$$\begin{cases} x_0 = x \in C \\ y_n = P_C(x_n - \lambda_n A x_n) \\ z_n = SP_C(x_n - \lambda_n A y_n) \\ C_n = \{z \in C : \|z_n - z\| \leq \|x_n - z\|\} \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\} \\ x_{n+1} = P_{C_n \cap Q_n} x \end{cases}$$

for every $n = 0, 1, 2, \dots$, where $\{\lambda_n\} \subset [a, b]$ for some $a, b \in (0, 1/k)$. Then the sequences $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ converge strongly to $P_{F(S) \cap VI(C, A)} x$.

Proof. It is obvious that C_n is closed and Q_n is closed and convex for every $n = 0, 1, 2, \dots$. As $C_n = \{z \in C : \|z_n - x_n\|^2 + 2\langle z_n - x_n, x_n - z \rangle \leq 0\}$, we also have C_n is convex for every $n = 0, 1, 2, \dots$. Put $t_n = P_C(x_n - \lambda_n A y_n)$ for every $n = 0, 1, 2, \dots$. Let $u \in F(S) \cap VI(C, A)$. From (3), monotonicity of A and $u \in VI(C, A)$, we have

$$\begin{aligned} \|t_n - u\|^2 &\leq \|x_n - \lambda_n A y_n - u\|^2 - \|x_n - \lambda_n A y_n - t_n\|^2 \\ &= \|x_n - u\|^2 - \|x_n - t_n\|^2 + 2\lambda_n \langle A y_n, u - t_n \rangle \\ &= \|x_n - u\|^2 - \|x_n - t_n\|^2 + 2\lambda_n (\langle A y_n - A u, u - y_n \rangle + \langle A u, u - y_n \rangle + \langle A y_n, y_n - t_n \rangle) \\ &\leq \|x_n - u\|^2 - \|x_n - t_n\|^2 + 2\lambda_n \langle A y_n, y_n - t_n \rangle \\ &= \|x_n - u\|^2 - \|x_n - y_n\|^2 - 2\langle x_n - y_n, y_n - t_n \rangle - \|y_n - t_n\|^2 + 2\lambda_n \langle A y_n, y_n - t_n \rangle \\ &= \|x_n - u\|^2 - \|x_n - y_n\|^2 - \|y_n - t_n\|^2 + 2\langle x_n - \lambda_n A y_n - y_n, t_n - y_n \rangle. \end{aligned}$$

Further, since $y_n = P_C(x_n - \lambda_n A x_n)$ and A is k -Lipschitz-continuous, we have

$$\begin{aligned} &\langle x_n - \lambda_n A y_n - y_n, t_n - y_n \rangle \\ &= \langle x_n - \lambda_n A x_n - y_n, t_n - y_n \rangle + \langle \lambda_n A x_n - \lambda_n A y_n, t_n - y_n \rangle \\ &\leq \langle \lambda_n A x_n - \lambda_n A y_n, t_n - y_n \rangle \\ &\leq \lambda_n k \|x_n - y_n\| \|t_n - y_n\|. \end{aligned}$$

So, we have

$$\begin{aligned} \|t_n - u\|^2 &\leq \|x_n - u\|^2 - \|x_n - y_n\|^2 - \|y_n - t_n\|^2 + 2\lambda_n k \|x_n - y_n\| \|t_n - y_n\| \\ &\leq \|x_n - u\|^2 - \|x_n - y_n\|^2 - \|y_n - t_n\|^2 + \lambda_n^2 k^2 \|x_n - y_n\|^2 + \|y_n - t_n\|^2 \\ &\leq \|x_n - u\|^2 + (\lambda_n^2 k^2 - 1) \|x_n - y_n\|^2 \\ &\leq \|x_n - u\|^2. \end{aligned} \tag{4}$$

Therefore from $z_n = St_n$ and $u = Su$, we have

$$\|z_n - u\| = \|St_n - Su\| \leq \|t_n - u\| \leq \|x_n - u\| \tag{5}$$

for every $n = 0, 1, 2, \dots$ and hence $u \in C_n$. So, $F(S) \cap VI(C, A) \subset C_n$ for every $n = 0, 1, 2, \dots$. Next, let us show by mathematical induction that $\{x_n\}$ is well-defined and $F(S) \cap VI(C, A) \subset C_n \cap Q_n$ for every $n = 0, 1, 2, \dots$. For $n = 0$ we have $Q_0 = C$. Hence we obtain $F(S) \cap VI(C, A) \subset C_0 \cap Q_0$. Suppose that x_k is given and $F(S) \cap VI(C, A) \subset C_k \cap Q_k$ for some $k \in N$. Since $F(S) \cap VI(C, A)$ is nonempty,

$C_k \cap Q_k$ is a nonempty closed convex subset of C . So, there exists a unique element $x_{k+1} \in C_k \cap Q_k$ such that $x_{k+1} = P_{C_k \cap Q_k} x$. It is also obvious that there holds $\langle x_{k+1} - z, x - x_{k+1} \rangle \geq 0$ for every $z \in C_k \cap Q_k$. Since $F(S) \cap VI(C, A) \subset C_k \cap Q_k$, we have $\langle x_{k+1} - z, x - x_{k+1} \rangle \geq 0$ for $z \in F(S) \cap VI(C, A)$ and hence $F(S) \cap VI(C, A) \subset Q_{k+1}$. Therefore, we obtain $F(S) \cap VI(C, A) \subset C_{k+1} \cap Q_{k+1}$.

Let $t_0 = P_{F(S) \cap VI(C, A)} x$. From $x_{n+1} = P_{C_n \cap Q_n} x$ and $t_0 \in F(S) \cap VI(C, A) \subset C_n \cap Q_n$, we have

$$\|x_{n+1} - x\| \leq \|t_0 - x\| \quad (6)$$

for every $n = 0, 1, 2, \dots$. Therefore, $\{x_n\}$ is bounded. We also have

$$\|z_n - u\| = \|St_n - Su\| \leq \|t_n - u\| \leq \|x_n - u\|$$

for some $u \in F(S) \cap VI(C, A)$. So, $\{z_n\}$ and $\{t_n\}$ are bounded. Since $x_{n+1} \in C_n \cap Q_n \subset Q_n$ and $x_n = P_{Q_n} x$, we have

$$\|x_n - x\| \leq \|x_{n+1} - x\|$$

for every $n = 0, 1, 2, \dots$. Therefore, there exists $c = \lim_{n \rightarrow \infty} \|x_n - x\|$. Since $x_n = P_{Q_n} x$ and $x_{n+1} \in Q_n$, we have

$$\begin{aligned} \|x_{n+1} - x_n\|^2 &= \|x_{n+1} - x\|^2 + \|x_n - x\|^2 + 2 \langle x_{n+1} - x, x - x_n \rangle \\ &= \|x_{n+1} - x\|^2 - \|x_n - x\|^2 - 2 \langle x_n - x_{n+1}, x - x_n \rangle \\ &\leq \|x_{n+1} - x\|^2 - \|x_n - x\|^2 \end{aligned}$$

for every $n = 0, 1, 2, \dots$. This implies that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0.$$

Since $x_{n+1} \in C_n$, we have $\|z_n - x_{n+1}\| \leq \|x_n - x_{n+1}\|$ and hence

$$\|x_n - z_n\| \leq \|x_n - x_{n+1}\| + \|x_{n+1} - z_n\| \leq 2 \|x_{n+1} - x_n\|$$

for every $n = 0, 1, 2, \dots$. From $\|x_{n+1} - x_n\| \rightarrow 0$, we have $\|x_n - z_n\| \rightarrow 0$.

For $u \in F(S) \cap VI(C, A)$, from (4) and (5) we obtain

$$\|z_n - u\|^2 \leq \|t_n - u\|^2 \leq \|x_n - u\|^2 + (\lambda_n^2 k^2 - 1) \|x_n - y_n\|^2.$$

Therefore, we have

$$\begin{aligned} \|x_n - y_n\|^2 &\leq \frac{1}{1 - \lambda_n^2 k^2} (\|x_n - u\|^2 - \|z_n - u\|^2) \\ &= \frac{1}{1 - \lambda_n^2 k^2} (\|x_n - u\| - \|z_n - u\|) (\|x_n - u\| + \|z_n - u\|) \\ &\leq \frac{1}{1 - \lambda_n^2 k^2} (\|x_n - u\| + \|z_n - u\|) \|x_n - z_n\|. \end{aligned}$$

Since $\|x_n - z_n\| \rightarrow 0$, we obtain $x_n - y_n \rightarrow 0$. From (4) and (5) we also have

$$\begin{aligned} \|z_n - u\|^2 &\leq \|t_n - u\|^2 \\ &\leq \|x_n - u\|^2 - \|x_n - y_n\|^2 - \|y_n - t_n\|^2 + 2\lambda_n k \|x_n - y_n\| \|t_n - y_n\| \\ &\leq \|x_n - u\|^2 - \|x_n - y_n\|^2 - \|y_n - t_n\|^2 + \|x_n - y_n\|^2 + \lambda_n^2 k^2 \|y_n - t_n\|^2 \\ &\leq \|x_n - u\|^2 + (\lambda_n^2 k^2 - 1) \|y_n - t_n\|^2. \end{aligned}$$

Therefore we have

$$\begin{aligned} \|t_n - y_n\|^2 &\leq \frac{1}{1 - \lambda_n^2 k^2} (\|x_n - u\|^2 - \|z_n - u\|^2) \\ &= \frac{1}{1 - \lambda_n^2 k^2} (\|x_n - u\| - \|z_n - u\|) (\|x_n - u\| + \|z_n - u\|) \\ &\leq \frac{1}{1 - \lambda_n^2 k^2} (\|x_n - u\| + \|z_n - u\|) \|x_n - z_n\|. \end{aligned}$$

Since $\|x_n - z_n\| \rightarrow 0$, we obtain $t_n - y_n \rightarrow 0$. Since A is k -Lipschitz-continuous, we have $Ay_n - At_n \rightarrow 0$. From $\|x_n - t_n\| \leq \|x_n - y_n\| + \|y_n - t_n\|$ we also have $x_n - t_n \rightarrow 0$. Since

$$\|t_n - St_n\| = \|t_n - z_n\| \leq \|t_n - x_n\| + \|x_n - z_n\|,$$

we have $\|t_n - St_n\| \rightarrow 0$.

As $\{x_n\}$ is bounded, there is a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $\{x_{n_i}\}$ converges weakly to some u . We can obtain that $u \in F(S) \cap VI(C, A)$. First, we show $u \in VI(C, A)$. Since $x_n - t_n \rightarrow 0$ and $x_n - y_n \rightarrow 0$, we have $\{t_{n_i}\} \rightarrow u$ and $\{y_{n_i}\} \rightarrow u$. Let

$$Tv = \begin{cases} Av + N_C v, & \text{if } v \in C, \\ \emptyset, & \text{if } v \notin C. \end{cases}$$

Then T is maximal monotone and $0 \in Tv$ if and only if $v \in VI(C, A)$; see [7]. Let $(v, w) \in G(T)$. Then, we have $w \in Tv = Av + N_C v$ and hence $w - Av \in N_C v$. So, we have $\langle v - t, w - Av \rangle \geq 0$ for all $t \in C$. On the other hand, from $t_n = P_C(x_n - \lambda_n Ay_n)$ and $v \in C$ we have

$$\langle x_n - \lambda_n Ay_n - t_n, t_n - v \rangle \geq 0$$

and hence

$$\left\langle v - t_n, \frac{t_n - x_n}{\lambda_n} + Ay_n \right\rangle \geq 0.$$

Therefore from $w - Av \in N_C v$ and $t_{n_i} \in C$, we have

$$\begin{aligned} \langle v - t_{n_i}, w \rangle &\geq \langle v - t_{n_i}, Av \rangle \\ &\geq \langle v - t_{n_i}, Av \rangle - \left\langle v - t_{n_i}, \frac{t_{n_i} - x_{n_i}}{\lambda_{n_i}} + Ay_{n_i} \right\rangle \\ &= \langle v - t_{n_i}, Av - At_{n_i} \rangle + \langle v - t_{n_i}, At_{n_i} - Ay_{n_i} \rangle - \left\langle v - t_{n_i}, \frac{t_{n_i} - x_{n_i}}{\lambda_{n_i}} \right\rangle \\ &\geq \langle v - t_{n_i}, At_{n_i} - Ay_{n_i} \rangle - \left\langle v - t_{n_i}, \frac{t_{n_i} - x_{n_i}}{\lambda_{n_i}} \right\rangle. \end{aligned}$$

Hence, we obtain $\langle v - u, w \rangle \geq 0$ as $i \rightarrow \infty$. Since T is maximal monotone, we have $u \in T^{-1}0$ and hence $u \in VI(C, A)$.

Let us show $u \in F(S)$. Assume $u \notin F(S)$. From Opial's condition, we have

$$\begin{aligned} \liminf_{i \rightarrow \infty} \|t_{n_i} - u\| &< \liminf_{i \rightarrow \infty} \|t_{n_i} - Su\| \\ &= \liminf_{i \rightarrow \infty} \|t_{n_i} - St_{n_i} + St_{n_i} - Su\| \\ &\leq \liminf_{i \rightarrow \infty} \|St_{n_i} - Su\| \\ &\leq \liminf_{i \rightarrow \infty} \|t_{n_i} - u\|. \end{aligned}$$

This is a contradiction. So, we obtain $u \in F(S)$. This implies $u \in F(S) \cap VI(C, A)$.

From $t_0 = P_{F(S) \cap VI(C, A)} x$, $u \in F(S) \cap VI(C, A)$ and (6), we have

$$\|t_0 - x\| \leq \|u - x\| \leq \liminf_{i \rightarrow \infty} \|x_{n_i} - x\| \leq \limsup_{i \rightarrow \infty} \|x_{n_i} - x\| \leq \|t_0 - x\|.$$

So, we obtain

$$\lim_{i \rightarrow \infty} \|x_{n_i} - x\| = \|u - x\|.$$

From $x_{n_i} - x \rightarrow u - x$ we have $x_{n_i} - x \rightarrow u - x$ and hence $x_{n_i} \rightarrow u$. Since $x_n \in P_{Q_n} x$ and $t_0 \in F(S) \cap VI(C, A) \subset C_n \cap Q_n \subset Q_n$, we have

$$-\|t_0 - x_{n_i}\|^2 = \langle t_0 - x_{n_i}, x_{n_i} - x \rangle + \langle t_0 - x_{n_i}, x - t_0 \rangle \geq \langle t_0 - x_{n_i}, x - t_0 \rangle.$$

As $i \rightarrow \infty$, we obtain $-\|t_0 - u\|^2 \geq \langle t_0 - u, x - t_0 \rangle \geq 0$ by $t_0 = P_{F(S) \cap VI(C, A)} x$ and $u \in F(S) \cap VI(C, A)$. Hence we have $u = t_0$. This implies that $x_n \rightarrow t_0$. It is easy to see $y_n \rightarrow t_0$, $z_n \rightarrow t_0$.

4 Applications.

Using Theorem 3.1, we prove some theorems in a real Hilbert space.

Theorem 4.1 *Let C be a closed convex subset of a real Hilbert space H . Let A be a monotone and k -Lipschitz-continuous mapping of C into H such that $VI(C, A)$ is nonempty. Let $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ be sequences generated by*

$$\begin{cases} x_0 = x \in C \\ y_n = P_C(x_n - \lambda_n A x_n) \\ z_n = P_C(x_n - \lambda_n A y_n) \\ C_n = \{z \in C : \|z_n - z\| \leq \|x_n - z\|\} \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\} \\ x_{n+1} = P_{C_n \cap Q_n} x \end{cases}$$

for every $n = 0, 1, 2, \dots$, where $\{\lambda_n\} \subset [a, b]$ for some $a, b \in (0, 1/k)$. Then the sequences $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ converge strongly to $P_{VI(C, A)}x$.

Proof. Putting $S = I$, by Theorem 3.1, we obtain the desired result.

Remark. See Iiduka, Takahashi and Toyoda [2] for the case when A is α -inverse-strongly-monotone.

Theorem 4.2 *Let C be a closed convex subset of a real Hilbert space H and S be a nonexpansive mapping of C into itself such that $F(S)$ is nonempty. Let $\{x_n\}$ and $\{y_n\}$ be sequences generated by*

$$\begin{cases} x_0 = x \in C \\ y_n = S x_n \\ C_n = \{z \in C : \|y_n - z\| \leq \|x_n - z\|\} \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\} \\ x_{n+1} = P_{C_n \cap Q_n} x \end{cases}$$

for every $n = 0, 1, 2, \dots$. Then the sequences $\{x_n\}$ and $\{y_n\}$ converge strongly to $P_{F(S)}x$.

Proof. Putting $A = 0$, by Theorem 3.1, we obtain the desired result.

Remark. See also Nakajo and Takahashi [5] for more general result.

Theorem 4.3 *Let H be a real Hilbert space. Let A be a monotone, k -Lipschitz-continuous mapping of H into itself and S be a nonexpansive mapping of H into itself such that $F(S) \cap A^{-1}0 \neq \emptyset$. Let $\{x_n\}$ and $\{y_n\}$ be sequences generated by*

$$\begin{cases} x_0 = x \in C \\ y_n = S(x_n - \lambda_n A(x_n - \lambda_n A x_n)) \\ C_n = \{z \in C : \|y_n - z\| \leq \|x_n - z\|\} \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\} \\ x_{n+1} = P_{C_n \cap Q_n} x \end{cases}$$

for every $n = 0, 1, 2, \dots$, where $\{\lambda_n\} \subset [a, b]$ for some $a, b \in (0, 1/k)$. Then the sequences $\{x_n\}$ and $\{y_n\}$ converge strongly to $P_{F(S) \cap A^{-1}0}x$.

Proof. We have $A^{-1}0 = VI(H, A)$ and $P_H = I$. By Theorem 3.1, we obtain the desired result.

Remark. Notice that $F(S) \cap A^{-1}0 \subset VI(F(S), A)$. See also Yamada [9] for the case when A is a strongly monotone and Lipschitz continuous mapping of a real Hilbert space H into itself and S is a nonexpansive mapping of H into itself.

Theorem 4.4 *Let H be a real Hilbert space. Let A be a monotone, k -Lipschitz-continuous mapping of H into itself and $B : H \rightarrow 2^H$ be a maximal monotone mapping such that $A^{-1}0 \cap B^{-1}0 \neq \emptyset$. Let J_r^B be the resolvent of B for each $r > 0$. Let $\{x_n\}$ and $\{y_n\}$ be sequences generated by*

$$\begin{cases} x_0 = x \in C \\ y_n = J_r^B(x_n - \lambda_n A(x_n - \lambda_n A x_n)) \\ C_n = \{z \in C : \|y_n - z\| \leq \|x_n - z\|\} \\ Q_n = \{z \in C : \langle x_n - z, x - x_n \rangle \geq 0\} \\ x_{n+1} = P_{C_n \cap Q_n} x \end{cases}$$

for every $n = 0, 1, 2, \dots$, where $\{\lambda_n\} \subset [a, b]$ for some $a, b \in (0, 1/k)$. Then the sequences $\{x_n\}$ and $\{y_n\}$ converge strongly to $P_{A^{-1}0 \cap B^{-1}0}x$.

Proof. We have $A^{-1}0 = VI(H, A)$ and $F(J_r^B) = B^{-1}0$. Putting $P_H = I$, by Theorem 3.1 we obtain the desired result.

References

- [1] F. E. Browder and W. V. Petryshyn, *Construction of fixed points of nonlinear mappings in Hilbert space*, J. Math. Anal. Appl. **20** (1967), 197-228.
- [2] H. Iiduka, W. Takahashi and M. Toyoda, *Approximation of solutions of variational inequalities for monotone mappings*, PanAmer. Math. J. **11** (2004), no. 2, 45-55.
- [3] G. M. Korpelevich, *The extragradient method for finding saddle points and other problems*, Matecon **12** (1976), 747-756.
- [4] F. Liu and M. Z. Nashed, *Regularization of nonlinear ill-posed variational inequalities and convergence rates*, Set-Valued Anal. **6** (1998), 313-344.
- [5] K. Nakajo and W. Takahashi, *Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups*, J. Math. Anal. Appl. **279** (2003), no. 2, 372-379. 463-478.
- [6] Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591-597.
- [7] R. T. Rockafellar, *On the maximality of sums of nonlinear monotone operators*, Trans. Amer. Math. Soc. **149** (1970), 75-88.
- [8] W. Takahashi, *Nonlinear Functional Analysis*, Yokohama Publishers, Yokohama, Japan, 2000.
- [9] I. Yamada, *The hybrid steepest-descent method for the variational inequality problem over the intersection of fixed-point sets of nonexpansive mappings*, in Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications (D. Butnariu, Y. Censor and S. Reich Eds.), Kluwer Academic Publishers, Dordrecht, Holland, 2001.